




FIG. 1. Weight in girls with 22q11 deletion syndrome, ages 0–36 months.




FIG. 2. Length in girls with 22q11 deletion syndrome, ages 0–36 months.




FIG. 3. Occipito-frontal head circumference in girls with 22q11 deletion syndrome, ages 0–36 months.




FIG. 7. Weight in females with 22q11 deletion syndrome, ages 2–20 years.






FIG. 8. Height in females with 22q11 deletion syndrome, ages 2–20 years.




FIG. 9. Occipito-frontal head circumference in females with 22q11 deletion syndrome, ages 2–20 years.




FIG. 10. BMI in females with 22q11 deletion syndrome, ages 2–20 years.




FIG. 4. Weight in boys with 22q11 deletion syndrome, ages 0–36 months.






FIG. 5. Length in boys with 22q11 deletion syndrome, ages 0–36 months.




FIG. 6. Occipito-frontal head circumference in boys with 22q11 deletion syndrome, ages 0–36 months.



FIG. 11. Weight in males with 22q11 deletion syndrome, ages 2–20 years.






FIG. 12. Height in males with 22q11 deletion syndrome, ages 2–20 years.






FIG. 13. Occipito-frontal head circumference in males with 22q11 deletion syndrome, ages 2–20 years.




FIG. 14. BMI in males with 22q11 deletion syndrome, ages 2–20 years.